

Mesenteric Vasodilation Mediated by Endothelial Anandamide Receptors

Jens A. Wagner, Károly Varga, Zoltán Járαι, George Kunos

Abstract—Cannabinoids, including the endogenous ligand anandamide (arachidonyl ethanolamide), elicit pronounced hypotension in rats via activation of peripherally located CB1 cannabinoid receptors, which have been also implicated in endotoxin (lipopolysaccharide [LPS])-induced hypotension. The present study was designed to test the role of vascular CB1 receptors in cannabinoid- and endotoxin-induced mesenteric vasodilation. In the isolated, buffer-perfused rat mesenteric arterial bed precontracted with phenylephrine, anandamide induced long-lasting (up to 60 minutes) dose-dependent vasodilation (ED_{50} : 79 ± 3 nmol; maximal relaxation: $77 \pm 2\%$), inhibited by 0.5 to 5.0 $\mu\text{mol/L}$ of the selective CB1 receptor antagonist SR141716A. Low doses of the calcium ionophore ionomycin also caused mesenteric vasodilation inhibited by SR141716A. The metabolically stable analogue R-methanandamide elicited mesenteric vasodilation (ED_{50} : 286 ± 29 nmol), whereas the potent synthetic CB1 receptor agonists WIN 55212-2 and HU-210 caused no change in vascular tone or only a minor dilator effect not affected by SR141716A, respectively. The endogenous ligand 2-arachidonyl glycerol caused no change in vascular tone, whereas Δ^9 -tetrahydrocannabinol and arachidonic acid caused mesenteric vasoconstriction. After endothelial denudation, the dilator response to anandamide was slightly reduced and was no longer inhibited by SR141716A. In preparations from LPS-pretreated rats, SR141716A alone caused a significant and prolonged increase in perfusion pressure, whereas it had no such effect in control preparations perfused in vitro with or without LPS or after endothelial denudation in preparations from rats pretreated with LPS. We conclude that anandamide-induced mesenteric vasodilation is mediated by an endothelially located SR141716A-sensitive “anandamide receptor” distinct from CB1 cannabinoid receptors and that activation of such receptors by an endocannabinoid, possibly anandamide, contributes to LPS-induced mesenteric vasodilation in vivo. (*Hypertension*. 1999;33[part II]:429-434.)

Key Words: vasodilation ■ cannabinoids ■ anandamide ■ endotoxin

Endogenous cannabinoids (or endocannabinoids) represent a novel class of lipid ligands that share receptor binding sites with plant-derived cannabinoids, such as Δ^9 -tetrahydrocannabinol (THC), and that mimic the neurobehavioral effects of the latter.¹ Two receptors have been identified by molecular cloning that can recognize cannabinoids with high affinity: the CB1 receptor is present primarily in the brain² but also in some peripheral tissues^{3,4} and the CB2 receptor is expressed by cells of the immune system.⁵ The mRNA of a splice variant of the CB1 receptor, CB1A, has also been identified.³ Two endocannabinoids have been characterized in some detail: arachidonyl ethanolamide, or anandamide,⁶ and 2-arachidonyl glyceride, or 2-AG.^{7,8} Neurons in the brain can synthesize anandamide⁹ and 2-AG,¹⁰ and both substances have been shown to influence central neural functions, such as long-term potentiation.^{10,11} Peripheral tissues also contain anandamide¹²⁻¹⁴ and 2-AG.^{7,15,16}

Plant-derived cannabinoids can produce cardiovascular effects, including hypotension,¹⁷ that also can be elicited by anandamide¹⁸⁻²⁰ and 2-AG.¹⁶ Although at first the hypoten-

sive effect of THC was thought to result from centrally mediated sympathoinhibition,¹⁷ more recent evidence implicates peripheral sites of action, such as receptors located on sympathetic nerve terminals,^{4,19,21} receptors located in vascular tissue, or both.^{16,20,22} In a recent study, the hypotensive potency of cannabinoid agonists, including anandamide, showed a strong positive correlation with the binding affinity of the same ligands to the brain cannabinoid receptor.²² Furthermore, the CB1 receptor antagonist SR141716A inhibited the neurobehavioral²³ and hypotensive effects of cannabinoids²² with similar potency. This suggests that cannabinoid-induced hypotension in anesthetized rats is mediated by CB1 receptors.²²

We recently reported that SR141716A antagonized the hypotension of hemorrhagic¹⁴ and endotoxic¹⁶ shock in rats, conditions that also induced circulating macrophages to generate anandamide^{14,16} and platelets to generate 2-AG.¹⁶ When isolated from animals in shock, these blood cells elicited SR141716A-sensitive hypotension in normal recipient rats, which suggested that macrophage- and platelet-

Received September 16, 1998; first decision October 2, 1998; revision accepted October 14, 1998.

From the Department of Pharmacology and Toxicology, Medical College of Virginia Commonwealth University, Richmond.

Correspondence to George Kunos, MD, PhD, Department of Pharmacology and Toxicology, MCV/VCU, PO Box 980613, Room 746, 410 North 12th St, Richmond, VA 23298. E-mail gkunos@hsc.vcu.edu

© 1999 American Heart Association, Inc.

Hypertension is available at <http://www.hypertensionaha.org>

derived endocannabinoids are likely contributors to shock-related hypotension.^{14,16} Although activated macrophages and platelets are known to adhere to the vascular wall, the mechanism by which mediators generated by these cells elicit relaxation of vascular smooth muscle and the potential additional role of the vascular endothelium have remained unclear. This is an important question because there is evidence that endocannabinoids may also be generated in the vascular endothelium.^{13,15} Furthermore, findings in the isolated perfused rat mesenteric arterial bed have led to the proposal that the elusive endothelium-derived hyperpolarizing factor may be an endocannabinoid,^{24,25} although this proposal has been challenged.^{26–28} The purpose of the present study was to test the role of CB1 receptors and of the vascular endothelium in cannabinoid- and endotoxin-induced mesenteric vasodilation.

Methods

Preparation of Rat Isolated Buffer-Perfused Mesenteric Arterial Bed

Male Sprague-Dawley rats (300 to 350 g) anesthetized with ether were laparotomized, and a second-order branch of the superior mesenteric artery was cannulated using a PE60 polyethylene cannula. The mesentery was isolated, separated from the intestines, and placed in a water-jacketed perfusion chamber maintained at 37°C.²⁹ The preparation was perfused at 2 mL/min with Krebs-Henseleit solution pre-gassed with 95% O₂/5% CO₂, with a peristaltic pump (Rainin). Perfusion pressure was monitored via a T tube inserted between the pump and the inflow cannula and connected to a pressure transducer (Abbott) and physiograph (Astromed). After a 30-minute equilibration period, the perfusion pressure was raised from 30 to 40 mm Hg to approximately 80 to 100 mm Hg by the inclusion of phenylephrine (15 μmol/L) in the medium. Once the perfusion pressure had stabilized, endothelium-dependent and -independent vasodilator responses were tested by bolus intra-arterial injections of acetylcholine and sodium nitroprusside, respectively. The magnitude of vasodilation (seen as a decrease in perfusion pressure) was expressed as percent relaxation of established tone. Agonists were injected as a 100-μL bolus into the artery over a period of 5 seconds, with their dose expressed as nmol of drug in 100 μL vehicle. Each preparation was tested with no more than 3 doses of an agonist. Antagonists were dissolved in the perfusion medium at the indicated concentrations (in μmol/L). In some experiments, rats received 15 mg/kg *Escherichia coli* (*E. coli*) lipopolysaccharide (LPS) IP 2 hours before removal of the mesenteric bed.

Endothelial Denudation

To achieve endothelial denudation, the preparation was perfused with distilled water in the absence of phenylephrine for 3 to 6 minutes. On resuming perfusion with Krebs-Henseleit buffer containing phenylephrine, perfusion pressure returned to the same level as before distilled water perfusion. Functional denudation was considered to be achieved when the maximal dilator response to acetylcholine was reduced to <20% of control or converted to a pressor response, whereas the maximal dilator response to sodium nitroprusside remained unchanged. Only those preparations that met these criteria were used for further testing.

Chemicals

SR141716A (N-[piperidin-1-yl]-5-[4-chlorophenyl]-1-[1,2-dichlorophenyl]-4-methyl-1H-pyrazole-3-carboxamide HCl) was from Sanofi Co; WIN 55212-2 ([R]-[+]-[2,3-dihydro-5-methyl-3-[[4-morpholinyl]methyl]pyrrolol[1,2,3-de]-1,4-benzoxazin-6-yl]-1[naphtaleny]l methanone mesylate), R(+)-methanandamide, and N^G-nitro-L-arginine methyl ester hydrochloride (L-NAME) were from RBI; THC (Δ⁹-tetrahydrocannabinol) and anandamide (arachidonyl




Figure 1. Dose-dependent mesenteric vasoconstrictor (A) or vasodilator (B) action of cannabinoid agonists and of arachidonic acid. Drugs were injected intra-arterially in a 100-μL bolus. Isolated, precontracted mesenteric arterial bed preparations were perfused as described in "Methods." Points and vertical lines represent mean ± SE from 4 to 6 separate experiments. The drugs tested were THC (●), arachidonic acid (■), anandamide (○), R-methanandamide (▽), HU-210 (◇), WIN 55212-2 (□), and 2-AG (△).

ethanolamide) were provided by Dr Billy R. Martin; HU-210 ([-]-11-OH-Δ⁹-THC) was from Dr Raphael Mechoulam; and 2-AG (2-arachidonyl-glyceride) was from Deva Biotech. Acetylcholine, sodium nitroprusside, phenylephrine, arachidonic acid, indomethacin, ionomycin, *E. coli* LPS (0127:B8), and phenylmethylsulfonyl fluoride (PMSF) were from Sigma Chemical Co. SR141716A, THC, anandamide, 2-AG, and HU-210 were dissolved in 1:1:18 emulphor:ethanol:saline. WIN 55212-2 was dissolved in 1:1:18 emulphor:DMSO:saline. Emulphor is a polyoxyethylated vegetable oil.

Statistical Analysis

The paired *t* test was used to compare agonist effects tested in the same preparations in the absence and in the presence of an antagonist. The statistical package of Tallarida and Murray³⁰ was used to determine agonist ED₅₀ values from graded dose-response curves.

Results

Pharmacological Characterization of Cannabinoid-Induced Mesenteric Vasodilation

Bolus doses of anandamide produced vasodilation that lasted 40 to 60 minutes (Figure 1). The ED₅₀ of anandamide was




Figure 2. Inhibition of the mesenteric vasodilator action of anandamide by SR141716A. Anandamide was administered as a bolus intra-arterial injection of the indicated dose in the absence (○) or presence (▼) of 0.5 $\mu\text{mol/L}$ or 5 $\mu\text{mol/L}$ (●) of SR141716A present in the perfusion medium. The antagonist was added to the medium 15 minutes before the injection of anandamide. Points and vertical bars represent mean \pm SE from 4 to 5 separate experiments.

79 \pm 3 nmol, the peak effect was 77 \pm 2%. Once perfusion pressure had returned to its preinjection level, a second injection of the same dose of anandamide produced a similar response. The metabolically stable analogue of anandamide, R-methanandamide,³¹ also caused dose-dependent relaxation, although it was somewhat less efficacious than anandamide (ED₅₀: 286 \pm 29 nmol, Figure 1). Because 2-AG was found to elicit a hypotensive response similar to that of anandamide,¹⁶ it was also tested in the mesenteric preparation. In bolus doses of 2.6 to 2600 nmol, 2-AG failed to alter perfusion pressure (Figure 1). In doses of 320 to 6400 nmol, THC caused only increases in perfusion pressure, without any evidence of

vasodilation (Figure 1). HU-210, a synthetic cannabinoid 3 orders of magnitude more potent than anandamide in causing profound hypotension in rats,²² caused only a very minor dilator effect at the ultrahigh bolus dose of 2.6 μmol , which was the same in the absence (18 \pm 2%, n=5) or presence of 1 $\mu\text{mol/L}$ SR141716A, a selective CB1 receptor inhibitor²³ (16 \pm 6%, n=4). Another potent hypotensive cannabinoid WIN 55212-2²² was also ineffective in eliciting mesenteric vasodilation in the dose range of 2 to 570 nmol (Figure 1).

Because these agonist effects did not fit the pharmacological profile of CB1 receptors, we tested whether the CB1 receptor antagonist SR141716A inhibited the vasodilator effect of anandamide. In the presence of 0.5 $\mu\text{mol/L}$ SR141716A, the anandamide dose-response curve was right-shifted by a factor of 8.3 (ED₅₀: 663 \pm 4 nmol, $P < 0.001$, Figure 2), which is comparable to SR141716A antagonism of CB1 receptor-mediated effects.²³ However, when the concentration of SR141716A was increased 10-fold (to 5 $\mu\text{mol/L}$), the degree of inhibition increased only 2-fold (ED₅₀ of anandamide: 1415 \pm 6 nmol, $P < 0.001$, Figure 2). This suggests that additional SR141716A-insensitive mechanisms also contribute to the vasodilator effect of anandamide. Because anandamide may act on the endothelium, on the vascular smooth muscle, or both, we tested the effects of anandamide in endothelium-denuded preparations.

Effects of Endothelial Denudation on Anandamide-Induced Mesenteric Vasodilation

Figure 3 illustrates the effects of denudation on the responses to acetylcholine and sodium nitroprusside. Anandamide retained its vasodilator action after denudation, although the effect was somewhat reduced ($P < 0.05$, Figure 4). However, this reduced effect was no longer influenced by SR141716A, as illustrated in Figure 4. This suggests that, although anandamide is capable of causing vasodilation by a direct effect on vascular smooth muscle, this effect is not mediated by an SR141716A-sensitive mechanism.




Figure 3. The effects of acetylcholine (ACh, 110 nmol), sodium nitroprusside (NP, 15 nmol), and anandamide (AN, 145 nmol) on vascular tone in an isolated, buffer-perfused mesenteric arterial bed preparation before (A) and after endothelial denudation (B). Endothelial denudation was achieved by a 4-minute perfusion with distilled water, as described in "Methods." Note that the effect of acetylcholine is converted from dilation to constriction, whereas the response to sodium nitroprusside remains unaffected and that of anandamide is only slightly reduced after denudation. PP indicates perfusion pressure.




Figure 4. The effect of endothelial denudation on the mesenteric vasodilator response to anandamide. The effect of 145 nmol anandamide was tested in control (left) and endothelium denuded (right) preparations, in the absence (open columns) or presence (shaded columns) of 5 $\mu\text{mol/L}$ SR141716A. * indicates a significant difference ($P < 0.05$) from corresponding values in the absence of antagonist. # indicates a significant difference from corresponding values in control preparations. Columns and vertical bars represent mean \pm SE from 5 to 7 experiments.

Is Anandamide-Induced Vasodilation Due to a Degradation Product of Anandamide?

Anandamide is rapidly degraded by an amidohydrolase,⁹ and the arachidonic acid thus released may account for noncannabinoid receptor-mediated effects.²⁸ However, this possibility could be ruled out in the mesenteric preparation. First, the serine protease inhibitor PMSF, which inhibits anandamide hydrolysis,³² did not prevent the dilator response to anandamide. In intact preparations, the dilator response to 29 nmol anandamide ($14 \pm 1\%$, $n=4$) was actually enhanced in the presence of 200 $\mu\text{mol/L}$ PMSF ($28 \pm 5\%$, $n=4$, $P < 0.05$), whereas in endothelium-denuded preparations a borderline potentiation was observed ($16 \pm 7\%$ versus $24 \pm 5\%$, $n=4$, $P=0.07$). Second, the metabolically stable analogue R-methanandamide (1.38 μmol) caused a $54 \pm 8\%$ dilation after denudation ($n=4$), which was similar to its effect in intact preparations ($51 \pm 4\%$, $n=5$, see Figure 1). Third, in bolus doses of 10 to 1000 nmol, arachidonic acid caused no dilation and only dose-dependent increases in perfusion pressure (Figure 1).

Role of the Endothelium in Endotoxin-Induced Mesenteric Vasodilation

In view of the postulated role of endocannabinoids and SR141716A-sensitive cannabinoid receptors in endotoxin-induced hypotension,¹⁶ we examined the possible involvement of the vascular endothelium in this effect. Perfusion of the phenylephrine precontracted mesenteric vasculature with 5.0 $\mu\text{mol/L}$ SR141716A did not influence vascular tone (1 ± 1 mm Hg, $n=7$). However, when the mesenteric arterial bed was isolated from rats pretreated with 15 mg/kg LPS, IP, 2 hours before they were killed, the inclusion of 0.5 or 5.0 $\mu\text{mol/L}$ SR141716A in the perfusion medium caused a sustained pressor response (Figure 5). After such preparations were first denuded, SR141716A when subsequently administered failed to affect perfusion pressure (Figure 5). In vitro perfusion of control preparations with LPS, 100 $\mu\text{g/mL}$ for 60 minutes ($n=5$), failed




Figure 5. The effect of SR141716A on perfusion pressure in mesenteric arterial preparations from control (open column) and LPS-pretreated (shaded columns) rats. Columns and vertical bars represent mean \pm SE from 4 to 6 separate preparations in each group. * indicates significant difference ($P < 0.05$) from value in control preparations.

to affect perfusion pressure. These findings suggest that LPS induces an SR141716A-sensitive endothelium-dependent dilatory response in the in situ blood-perfused mesenteric vasculature.

The biosynthesis of anandamide is calcium dependent, and ionomycin has been used to induce anandamide synthesis in various cell types.^{9,12,14} In micromolar doses, ionomycin was found to elicit irreversible mesenteric vasoconstriction, probably due to a direct effect on vascular smooth muscle (results not shown). However, when used at extremely low doses of 10 to 100 pmol per 100 μL bolus, ionomycin elicited reversible and reproducible dilator responses, which were significantly inhibited in the presence of 5 $\mu\text{mol/L}$ SR141716A (Figure 6), similar to the observations of Randall et al.²⁴ In all experiments with ionomycin, the perfusion buffer contained 100 $\mu\text{mol/L}$ L-NAME and 10 $\mu\text{mol/L}$ indomethacin to block nitric oxide synthase and cyclooxygenase, respectively. Endothelial denudation reduced the dilator response to ionomycin, and the residual response was no longer affected by SR141716A (Figure 6).

Discussion

The present findings indicate that THC and the highly potent synthetic cannabinoids HU-210 and WIN 55212-2, which produce profound and long-lasting hypotension in anesthetized rats via activation of CB1 cannabinoid receptors,²² are devoid of mesenteric vasodilator activity or, in the case of HU-210, caused a minor dilator response not mediated by CB1 receptors. On the other hand, the endogenous ligand anandamide, which causes relatively modest and brief (<10 minutes) hypotension in anesthetized rats in vivo,^{18,19,22} elicits near-maximal and long-lasting (>30 minutes) mesenteric vasodilation. The metabolically stable analogue R-methanandamide has a similar effect. Another endogenous ligand, 2-AG, which can also elicit hypotension in anesthetized rats,¹⁶ has no mesenteric vasodilator activity. These findings indicate that, first, the mesenteric vasculature does not significantly contribute to the profound hypotension




Figure 6. The endothelium-dependent vasodilator effect of ionomycin is antagonized by SR141716A. The effect of a bolus injection of 100 pmol ionomycin was tested in control (left) and endothelium-denuded (right) preparations, in the absence (open columns) or presence (gray columns) of 5 $\mu\text{mol/L}$ SR141716A. In both cases, the perfusion buffer contained 100 $\mu\text{mol/L}$ L-NAME and 10 $\mu\text{mol/L}$ indomethacin. * indicates a significant difference ($P < 0.05$) from corresponding value in the absence of SR141716A; # indicates a significant difference ($P < 0.01$) from corresponding values in control preparations. Columns and vertical bars represent mean \pm SE from 4 to 5 experiments.

caused *in vivo* by plant-derived or synthetic cannabinoids and that, second, the receptors mediating the mesenteric vasodilator response to anandamide and R-methanandamide are distinct from CB1 receptors. These receptors should preferably be called anandamide receptors rather than cannabinoid receptors, to reflect their insensitivity to compounds derived from or related to cannabis.

The role of CB1 receptors in cannabinoid-induced hypotension was indicated by earlier findings with agonists and antagonists.²² The hypotensive potency of a series of agonists, including those tested in the present study, displayed a strong positive correlation with their analgesic potency or their binding K_d to the brain cannabinoid receptor.²² Furthermore, the inhibitory potency of the highly selective CB1 receptor antagonist SR141716A²³ was similar for cannabinoid-induced hypotension³³ and neurobehavioral effects.²³ Although SR141716A did inhibit the mesenteric vasodilator response to anandamide, the inhibition was unusual in that a 10-fold increase in the concentration of the antagonist resulted in a much smaller than expected increase in inhibition. Such "ceiling" effects, which result in anomalous Schild plots, can arise if more than one mechanism is involved in the action of the agonist, only one of which is sensitive to the antagonist. Indeed, SR141716A failed to inhibit anandamide-induced vasodilation in endothelium-denuded preparations. This suggests that mesenteric vasodilation by anandamide has 2 components: one mediated by a SR141716A-sensitive non-CB1 receptor site located on the endothelium and the other by an SR141716A-resistant direct action on vascular smooth muscle. The relative contribution of these 2 components to the net response to anandamide may depend on the species and tissue used and on the condition of the endothelium, which may explain conflicting reports in the

literature about the ability^{24,25} or inability^{26–28} of SR141716A to block the vasorelaxant effect of anandamide.

The SR141716A-resistant effect of anandamide cannot be attributed to arachidonic acid or its metabolites because in endothelium-denuded preparations PMSF, which prevents the degradation of anandamide, did not antagonize anandamide-induced vasodilation, and the metabolically stable R-methanandamide retained its vasorelaxant property. It is unclear whether the SR141716A-resistant effect of anandamide is unrelated to cannabinoid receptors, such as a direct opening of potassium channels, or whether it is mediated by an as-yet-unidentified cannabinoid receptor in vascular smooth muscle cells. However, CB2 receptors are unlikely to be involved because (1) anandamide binds to but does not activate CB2 receptors,³⁴ (2) the potent CB2 agonist WIN 55212-2³⁵ failed to cause mesenteric vasorelaxation, and (3) SR141716A, which has a K_d of 700 nmol/L at CB2 receptors,²³ should have inhibited the response at the concentration used.

In mesenteric preparations from LPS-pretreated rats, the endothelium-dependent pressor response to SR141716A indicates a role for the endothelium in endotoxin-induced mesenteric vasodilation. This finding, together with the observation that low doses of a calcium ionophore elicit endothelium-dependent SR141716A-sensitive mesenteric vasodilation, suggests that an endocannabinoid may be released from the endothelium of mesenteric arterioles. Indeed, anandamide has been identified in endothelial cells in the renal vasculature,¹³ and 2-AG was found to be present in endothelial cells from a human umbilical vein.¹⁵ However, in the rat mesenteric arteriolar bed, 2-AG does not cause dilation, and the SR141716A-sensitive component of the dilatory response to anandamide is endothelium dependent. Therefore, if anandamide is the cannabinoid released from mesenteric endothelial cells by ionomycin or LPS, its primary site of action must also be the endothelium, which implies its luminal release. Because of the presence of cholinacetyltransferase in vascular endothelial cells, a similar mechanism has been proposed for the endothelium-dependent vasodilatory effect of acetylcholine.³⁶ However, in contrast to the effect of acetylcholine, the putative endothelial mediator of SR141716A-sensitive vasodilation is unlikely to be nitric oxide, prostacyclin, or both, because the effect of ionomycin was observed in the presence of L-NAME and indomethacin.

An alternative or additional source of anandamide in the LPS-pretreated preparations may be circulating macrophages. LPS treatment has been shown to induce the production of anandamide by macrophages,¹⁶ and activated macrophages may actually become incorporated into the endothelium in which they can form tight junctions with endothelial cells.³⁷ Anandamide released from such resident macrophages may act in a juxtacrine manner to activate endothelial cannabinoid receptors. That macrophages may be required for the pressor effect of SR141716A in the LPS-treated preparations is suggested by the inability of LPS to induce mesenteric vasodilation *in vitro*, in a buffer-perfused preparation. Although LPS can directly activate the vascular endothelium, this mechanism is far less potent than the indirect pathway of endothelial activation dependent on circulating macrophages.³⁸

In summary, the present findings indicate that anandamide induces mesenteric vasodilation in the rat via a unique receptor located on endothelial cells, which is not activated by other cannabinoids but can be inhibited by the CB1-selective antagonist SR141716A. This endothelial receptor is thus distinct from CB1 receptors that mediate the profound hypotensive effect of synthetic cannabinoids *in vivo*. Activation of endothelial anandamide receptors may contribute to mesenteric vasodilation in endotoxic shock.

Acknowledgments

This study was supported by NIH Grants HL-49938 and HL-55297 to Dr Kunos and Grant VA-96-GS7 from the American Heart Association Virginia Affiliate to Dr Varga. Dr Wagner was supported by a fellowship from the Deutsche Forschungsgemeinschaft, and Dr J arai was supported by a fellowship from Sanofi Research Inc (Montpellier, France). We thank Dr Billy R. Martin (Richmond, VA) for helpful discussions and Dr Raphael Mechoulam (Jerusalem, Israel) for the gift of HU-210 used in this study.

References

- Mechoulam R, Shabat SB, Hanus L, Fried E, Bayewitch M, Vogel Z. Endogenous cannabinoid ligands. *Adv Exp Biol Med*. 1996;402:95–101.
- Matsuda LA, Lolait SJ, Brownstein MJ, Young AC, Bonner TI. Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature*. 1990;346:561–564.
- Shire D, Carillon C, Kaghad M, Calandra B, Rinaldi-Carmona M, Le Fur G, Ferrara P. An amino-terminal variant of the central cannabinoid receptor resulting from alternative splicing. *J Biol Chem*. 1995;270:3726–3731.
- Ishac E, JN, Jiang L, Lake KD, Varga K, Abood MA, Kunos G. Inhibition of exocytotic noradrenaline release by presynaptic cannabinoid CB1 receptors on peripheral sympathetic nerves. *Br J Pharmacol*. 1996;118:2023–2028.
- Munro S, Thomas KL, Abu-Shaar M. Molecular characterization of a peripheral receptor for cannabinoids. *Nature*. 1993;365:61–65.
- Devane WA, Hanus L, Breuer A, Pertwee RG, Stevenson LA, Griffin G, Gibson D, Mandelbaum A, Etinger A, Mechoulam R. Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science*. 1992;258:1946–1949.
- Mechoulam R, Ben-Shabat S, Hanus L, Ligumsky M, Kaminsky NE, Schatz AR, Gopher A, Almog S, Martin BR, Compton DR, Pertwee RG, Griffin G, Bayewitch M, Barg J, Vogel Z. Identification of an endogenous 2-monoglyceride, present in canine gut, that binds to cannabinoid receptors. *Biochem Pharmacol*. 1995;50:83–90.
- Sugiura T, Kondo S, Sukagawa A, Nakane S, Shinoda A, Itoh K, Yamashita A, Waku K. 2-Arachidonoyl-glycerol: a possible endogenous cannabinoid receptor ligand in brain. *Biochem Biophys Res Commun*. 1995;215:89–97.
- Di Marzo V, Fontana A, Cadas H, Schinelli S, Cimino G, Schwartz J-C, Piomelli D. Formation and inactivation of endogenous cannabinoid anandamide in central neurons. *Nature*. 1994;372:686–691.
- Stella N, Schweitzer P, Piomelli D. A second endogenous cannabinoid that modulates long-term potentiation. *Nature*. 1997;388:773–778.
- Collin C, Devane WA, Dahl D, Lee CJ, Axelrod J, Alkon DL. Long-term synaptic transformation of hippocampal CA1 gamma-aminobutyric acid synapses and the effect of anandamide. *Proc Natl Acad Sci U S A*. 1995;92:10167–10171.
- Di Marzo V, De Petrocellis L, Sepe N, Buono A. Biosynthesis of anandamide and related acylethanolamides in mouse J774 macrophages and N₁₈ neuroblastoma cells. *Biochem J*. 1996;316:977–984.
- Deutsch DG, Goligorsky MS, Schmid PC, Krebsbach RJ, Schmid HH, Das SK, Dey SK, Arreza G, Thorup C, Stefano G, Moore LC. Production and physiological actions of anandamide in the vasculature of the rat kidney. *J Clin Invest*. 1997;100:1538–1546.
- Wagner JA, Varga K, Ellis EF, Rzigalinski BA, Martin BR, Kunos G. Activation of peripheral CB1 cannabinoid receptors in haemorrhagic shock. *Nature*. 1997;390:518–521.
- Sugiura T, Kodaka T, Nakane S, Kishimoto S, Kondo S, Waku K. Detection of an endogenous cannabimimetic molecule, 2-arachidonoylglycerol, and cannabinoid CB1 receptor mRNA in human vascular cells: is 2-arachidonoylglycerol a possible vasomodulator? *Biochem Biophys Res Commun*. 1998;243:838–843.
- Varga K, Wagner JA, Bridgen DT, Kunos G. Platelet- and macrophage-derived endogenous cannabinoids are involved in endotoxin-induced hypotension. *FASEB J*. 1998;12:1035–1044.
- Vollmer RR, Cavero I, Ertel RJ, Solomon TA, Buckley JP. Role of the central autonomic nervous system in the hypotension and bradycardia induced by (-)-Δ⁹-trans-tetrahydrocannabinol. *J Pharm Pharmacol*. 1974;26:186–198.
- Varga K, Lake K, Martin BR, Kunos G. Novel antagonist implicates the CB1 cannabinoid receptor in the hypotensive action of anandamide. *Eur J Pharmacol*. 1995;278:279–283.
- Varga K, Lake KD, Huangfu D, Guyenet P, Kunos G. Mechanism of the hypotensive action of anandamide in anesthetized rats. *Hypertension*. 1996;28:682–688.
- Calignano A, La Rana G, Beltramo M, Makryannis A, Piomelli D. Potentiation of anandamide hypotension by the transport inhibitor, AM404. *Eur J Pharmacol*. 1997;337:R1–R2.
- Malinowska B, Godlewski G, Bucher B, Schlicker E. Cannabinoid CB1 receptor-mediated inhibition of the neurogenic vasopressor response in the pithed rat. *Naunyn Schmiedebergs Arch Pharmacol*. 1997;281:1030–1037.
- Lake KD, Compton DR, Varga K, Martin BR, Kunos G. Cannabinoid-induced hypotension and bradycardia in rats is mediated by CB₁-like cannabinoid receptors. *J Pharmacol Exp Ther*. 1997;281:1030–1037.
- Rinaldi-Carmona M, Barth F, H eaulme M, Shire D, Calandra B, Congy C, Martinez S, Marani J, N eliat G, Caput D, Ferrara P, Soubri e P, Breli ere JC, Le Fur G. SR141716A, a potent and selective antagonist of the brain cannabinoid receptor. *FEBS Lett*. 1994;350:240–244.
- Randall MD, Alexander SPH, Bennett T, Boyd EA, Fry JR, Gardiner SM, Kemp PA, McCulloch AI, Kendall DA. An endogenous cannabinoid as an endothelium-derived vasorelaxant. *Biochem Biophys Res Commun*. 1996;229:114–120.
- White R, Hiley CR. A comparison of EDHF-mediated and anandamide-induced relaxations in the rat isolated mesenteric artery. *Br J Pharmacol*. 1997;122:1573–1584.
- Plane F, Holland M, Waldron GJ, Garland CJ, Boyle JP. Evidence that anandamide and EDHF act via different mechanisms in rat isolated mesenteric arteries. *Br J Pharmacol*. 1997;121:1509–1511.
- Chataigneau T, F el etou M, Thollon C, Villeneuve N, Vilaine J-P, Duhault J, Vanhoutte PM. Cannabinoid CB₁ receptor and endothelium-dependent hyperpolarization in guinea-pig carotid, rat mesenteric and porcine coronary arteries. *Br J Pharmacol*. 1998;123:968–974.
- Pratt PF, Hillard CJ, Edgemond WS, Campbell WB. N-arachidonylethanolamide relaxation of bovine coronary artery is not mediated by CB1 cannabinoid receptor. *Am J Physiol*. 1998;274:H375–H381.
- McGregor DD. The effect of sympathetic nerve stimulation on vasoconstrictor responses in perfused mesenteric blood vessels of the rat. *J Physiol (Lond)*. 1965;177:21–30.
- Tallarida RJ, Murray RB. *Manual of Pharmacologic Calculations with Computer Programs*. 2nd edition. New York, NY: Springer; 1987.
- Abadji V, Lin S, Taha G, Griffin G, Stevenson LA, Pertwee RG, Makryannis A. (R)-methanandamide: a chiral novel anandamide possessing higher potency and metabolic stability. *J Med Chem*. 1994;37:1889–1893.
- Deutsch DG, Chin SA. Enzymatic synthesis and degradation of anandamide, a cannabinoid receptor agonist. *Biochem Pharmacol*. 1993;46:791–796.
- Lake KD, Martin BR, Kunos G, Varga K. Cardiovascular effects of anandamide in anesthetized and conscious normotensive and hypertensive rats. *Hypertension*. 1997;29:1204–1210.
- Schatz AR, Lee M, Condie RB, Pulaski JT, Kaminski NE. Cannabinoid receptors CB1 and CB2: a characterization of expression and adenylate cyclase modulation within the immune system. *Toxicol Appl Pharmacol*. 1997;142:278–287.
- Showalter VM, Compton DR, Martin BR, Abood MA. Evaluation of binding in a transfected cell line expressing a peripheral cannabinoid receptor (CB2): identification of cannabinoid receptor subtype selective ligands. *J Pharmacol Exp Ther*. 1996;278:989–999.
- Parnavelas JG, Kelly W, Burnstock G. Ultrastructural localization of choline acetyltransferase in vascular endothelial cells in rat brain. *Nature*. 1985;316:724–725.
- Jara PI, Boric MP, Saez JC. Leukocytes express connexin 43 after activation with lipopolysaccharide and appear to form gap junctions with endothelial cells after ischemia-reperfusion. *Proc Natl Acad Sci U S A*. 1995;92:7011–7015.
- Pugin J, Ulevitch RJ, Tobias PS. A critical role for monocytes and CD14 in endotoxin-induced endothelial cell activation. *J Exp Med*. 1993;178:2193–2200.